

## ITLS Advanced Pre-Test

- 1 . A 34 year old man is injured in a motor vehicle collision. Physical examination is normal except for diffuse abdominal tenderness. Capillary refill is normal. Vital signs obtained en route to the hospital are BP 100/60; pulse, 120/min; respirations 28/min. How should this patient be managed in the field?
  - A. intravenous (I.V.) of 0.9% normal saline (NS) at "keep open" rate
  - B. I.V. of RL at 20 ml/kg bolus
  - C. pneumatic antishock trousers (PASG); inflate all three compartments
  - D. insert an oropharyngeal airway and assist ventilations
- 2 . What is Sellick's maneuver?
  - A. a method allowing the rescuer to hold a mask on the face with both hands
  - B. a system used to calculate minute volume
  - C. another name for Mallampati
  - D. posteriorly directed pressure applied to the cricoid cartilage
- 3 . Which one of the following viruses is there currently an effective vaccine?
  - A. delta hepatitis
  - B. hepatitis B
  - C. hepatitis C
  - D. human immunodeficiency virus
- 4 . A 16 year old girl is rescued from a burning house. She has 25% partial thickness burns, and the burned areas are hot to the touch. What is the appropriate treatment?
  - A. apply ice to burned areas until cool to touch
  - B. apply clean water to burned areas for up to 1-2 minutes
  - C. apply iced water to burned areas until cool to touch
  - D. do not apply anything to burned areas other than clean sheets
- 5 . Proper immobilization of a forearm (radius and ulna) fracture involves splinting which of the following?
  - A. elbow and fracture site
  - B. fracture site only
  - C. wrist and fracture site
  - D. wrist, elbow, and fracture site
- 6 . What treatment is NOT indicated in the routine management of the patient with a head injury?
  - A. administration of 100% oxygen
  - B. fluid resuscitation to a BP of 110-120 systolic if the patient is hypotensive
  - C. hyperventilation
  - D. stabilization of the cervical spine
- 7 . Which of the following sets of vital signs is most compatible with a diagnosis of isolated head injury with increasing intracranial pressure?
  - A. BP 170/100, pulse 50/min
  - B. BP 80/60, pulse 130/min
  - C. BP 80/60, pulse 50/min
  - D. BP 170/100, pulse 130/min
- 8 . Which of the following trauma situations would be considered "load and go"?
  - A. awake and alert patient who has blood coming out of one ear
  - B. a patient who had brief loss of consciousness but is now awake
  - C. patient with abdominal tenderness, clammy skin, blood pressure 100/70, and pulse 130/min
  - D. patient with tender, deformed lower leg

## ITLS Advanced Pre-Test

- 9 . Which one of the following should be performed, at the scene of a "load & go", prior to moving the trauma patient to the ambulance?
- A. apply traction splint for femur fracture
  - B. decompress tension pneumothorax
  - C. initiate intravenous line
  - D. obtain vital signs
- 10 . Which one of the following is a reason to interrupt the initial assessment?
- A. cardiac arrest
  - B. multiple open (compound) fractures
  - C. severe head injury with brain tissue visible
  - D. severe shock
- 11 . A 24 year old woman is found lying on the sidewalk after jumping from a fourth story window. She is hypotensive, diaphoretic, tachycardic, and unconscious. Injuries include open skull fracture, pelvis fractures, and flail chest. What should be your goal for your scene?
- A. 5 minutes or less
  - B. 5 to 10 minutes
  - C. 10 to 15 minutes
  - D. as long as necessary to assess and stabilize the patient
- 12 . Which of the following injuries would change an otherwise stable trauma patient's category from "stable" to "load and go"?
- A. bilateral clavicle fractures
  - B. bilateral femur fractures
  - C. bilateral humerus fractures
  - D. bilateral tibia fractures
- 13 . What is most commonly injured, during a fall from a height, of an infant?
- A. head
  - B. chest
  - C. abdomen
  - D. extremities
- 14 . What site is first choice for pediatric intraosseous infusion?
- A. proximal tibia
  - B. distal humerus
  - C. proximal femur
  - D. distal fibula
- 15 . Which of the following will generally suffer the LEAST structural damage from a gunshot wound from a rifle?
- A. spleen
  - B. kidney
  - C. liver
  - D. lung
- 16 . A 34 year old man has a gunshot wound to the right groin area. Arterial bleeding, which cannot be controlled with direct pressure, is coming from the wound. The patient appears confused, diaphoretic, and has weak peripheral pulses. What is the appropriate fluid resuscitation for this patient?
- A. I.V. at "keep open" rate
  - B. Apply a hemostatic agent and gain IV access given enough fluid to maintain peripheral pulses
  - C. I.V. wide open rate; give at least two liters, then reassess patient
  - D. no I.V. should be started in this situation

## ITLS Advanced Pre-Test

- 17 . Which one of the following is typically associated with, post-traumatic hemorrhage, EARLY shock?
- A. ventricular dysrhythmias
  - B. hypotension
  - C. loss of 30% to 45% of blood volume
  - D. narrowed pulse pressure
- 18 . Among the following, what is the most common cause of preventable trauma death in the injured adult patient?
- A. airway obstruction
  - B. cardiac tamponade
  - C. hemorrhagic shock
  - D. spinal injury
- 19 . In which of the following situations should an emergency rescue be used?
- A. leaking antifreeze from radiator
  - B. paralyzed patient
  - C. presence of toxic fumes
  - D. pregnant patient
- 20 . Which of the following is most typical of early, neurogenic shock?
- A. increased pulse, clammy skin
  - B. increased pulse, warm and dry skin
  - C. decreased pulse, clammy skin
  - D. decreased pulse, warm and dry skin
- 21 . A 23 year old man is injured in a motorcycle collision. The patient appears disoriented and grossly intoxicated. There is a large laceration on his scalp which is actively bleeding. The patient refuses treatment and threatens to call his lawyer if anyone touches him. What should you do?
- A. allow the patient to phone his lawyer
  - B. have the patient sign a release form, then let him go
  - C. have the patient placed under protective custody then treat and transport the patient using restraints if necessary
  - D. wait until the patient passes out from his head injury or bleeding then transport
- 22 . Which of the following is an acceptable location to insert a needle when decompressing a tension pneumothorax?
- A. directly under the bottom of the second rib, midclavicular line
  - B. directly under the bottom of the third rib, midclavicular line
  - C. directly over the top of the third rib, midaxillary line
  - D. directly over the top of the third rib, midclavicular line
- 23 . A 54 year old man is involved in a motor vehicle collision. The steering wheel is bent. During your initial assessment you note his skin is pale and his radial pulses are present. Breath sounds are clear, heart tones are not muffled. Which one of the following is most consistent with these
- A. cardiac contusion
  - B. traumatic aortic rupture
  - C. flail chest
  - D. tension pneumothorax
- 24 . What is the most common cause of cardiopulmonary arrest in the trauma patient?
- A. brain injury
  - B. hypoxemia
  - C. myocardial contusion
  - D. ventricular arrhythmia

## ITLS Advanced Pre-Test

- 25 . A 49-year-old man is involved in a motor vehicle collision. First Responders are doing CPR. Findings include a distended abdomen and obviously deformed pelvis and quick look on the monitor shows asystole. Which of the following is the most appropriate act?
- A. Establish IV access and administer a 20mL/kg bolus
  - B. Establish IV access and administer a 1 liter bolus
  - C. Establish IV access and administer a 2-4 liter bolus
  - D. Resuscitative efforts should not be started and the patient pronounced dead
- 26 . Which of the following statements concerning treatment of shock in the pregnant, burn patient is TRUE?
- A. oxygen should be used sparingly so as to avoid oxygen toxicity to the fetus
  - B. pressor agents such as dopamine should be used to improve circulation to the fetus
  - C. volume replacement should be given earlier and in larger amounts to the pregnant, burn patient
  - D. volume replacement should be given more slowly so as to avoid fluid overloading the fetus
- 27 . Which area of the spine is most susceptible to injury in a rear-impact motor vehicle crash?
- A. cervical
  - B. thoracic
  - C. lumbar
  - D. sacral-coccygeal
- 28 . An unrestrained 18 y/o male on the way to a post-graduation party leaves the road, bounces through a ditch and hits a tree. You find him behind the bent steering wheel, unconscious, cool, pale and clammy, blue around the lips with labored respirations of 30 and shallow, thready radial pulses of about 120, distended neck veins, tracheal deviation to the right, an asymmetrical chest with absent breath sounds on the left. You assume he has a
- A. Cardiac tamponade
  - B. Tension pneumothorax
  - C. Massive hemothorax
  - D. Simple pneumothorax
- 29 . Which one of the following mnemonics can be used to help predict which patients might potentially have difficult laryngoscopy and intubation?
- A. MMAP
  - B. IPPV
  - C. RSI
  - D. BOOTS
- 30 . An unrestrained 18 y/o male on the way to a post-graduation party leaves the road, bounces through a ditch and hits a tree. You find him behind the bent steering wheel, unconscious, cool, pale and clammy, with labored respirations of 30 and shallow, thready radial pulses of about 120, flat neck veins, trachea midline, an asymmetrical chest with absent breath sounds on the left. You assume he has a
- A. Cardiac tamponade
  - B. Tension pneumothorax
  - C. Massive hemothorax
  - D. Simple pneumothorax

## ITLS Advanced Pre-Test

- 31 .** A 23 y/o female has won the Let's-see-who-can-lean-the-farthest-backward-over-the-second-story-balcony-railing contest. You arrive to find her boyfriend standing over her, holding two beers, as she lies on the grass under the balcony. She opens her eyes to voice, her skin is normal in color, respirations about 16 and unlabored, pulse 54 and a little weak at the wrist, with no external bleeding. She has flat neck veins, a normal chest and abdomen and a stable pelvis. 1st responders who arrived just before you tell you that her pulse ox reading is 94 and her BP is 74/30. If this is true,
- A.** Hypovolemic shock
  - B.** Relative hypovolemic (high-space) shock
  - C.** Mechanical (obstructive) shock
  - D.** Cardiogenic shock
- 32 .** A 23 y/o female has won the Let's-see-who-can-lean-the-farthest-backward-over-the-second-story-balcony-railing contest. You arrive to find her boyfriend standing over her, holding two beers, as she lies on the grass under the balcony. She opens her eyes to voice, her skin is cool, clammy and ashen, respirations are rapid and shallow, radial pulses are too rapid to count and thready. There is no external bleeding. She has flat neck veins, a normal chest and abdomen and pelvis. Her boyfriend tells you that she has some kind of heart problem but he doesn't know what. Placed on the monitor, she shows a wide-complex tachycardia of about 280. What kind of shock is she suffering?
- A.** Hypovolemic shock
  - B.** Relative hypovolemic (high-space) shock
  - C.** Mechanical (obstructive) shock
  - D.** Cardiogenic shock
- 33 .** Which of the following would be the most compelling reason to intubate a patient immediately?
- A.** Snoring respirations
  - B.** Gurgling respirations
  - C.** Inability to ventilate (achieve chest rise)
  - D.** Traumatic arrest